

Analyses of Commercial Parrotfish Landings in the U.S. Caribbean
NOAA National Marine Fisheries Service
Southeast Regional Office
March 8, 2012

Introduction

Parrotfish are both a culturally and ecologically significant species in the Caribbean. They are harvested throughout the U.S. Caribbean with the majority of commercial landings coming from St. Croix. Parrotfish are herbivorous grazers that remove algae, which enhances settlement and survival of coral recruits (Brock 1979; Mumby 2006; Burkepille and Hay 2010). Their ecological role has become more relevant in the past 30 years due to the Caribbean-wide decline of longspine urchin (*Diadema antillarum*) in the 1980s (Mumby 2006).

In recent years, the Caribbean Fishery Management Council (CFMC) has approved several regulations to directly or indirectly reduce parrotfish harvest. In November 2005, a prohibition of gill and trammel nets was implemented in federal waters (50 CFR Parts 600 and 622). A compatible regulation was implemented in state waters of the U.S. Virgin Islands (USVI) in July 2006 (V.I.C. Title 12, Chapter 9A, 321-1), but, the Commonwealth of Puerto Rico did not enact similar regulations in state waters. Amendment 5 to the CFMC's Reef Fish Fishery Management Plan (FMP) was implemented on January 30, 2012, and prohibited the harvest of blue (*Scarus coeruleus*), midnight (*Scarus coelestinus*), and rainbow (*Scarus guacamaia*) parrotfish. These are the three largest parrotfish species that occur in the U.S. Caribbean and are considered to be the most vulnerable to overharvest because of their large body size, high susceptibility to spear gear and fish traps, and relatively low resilience.

The Magnuson-Stevens Reauthorization Act of 2006 required regional fishery management councils under federal management to implement annual catch limits (ACLs) and accountability measures (AMs) for all stocks undergoing overfishing by 2010. The CFMC implemented ACLs for parrotfish in Amendment 5 to the CFMC's Reef Fish FMP (CFMC 2010). ACLs were implemented for each island based on 85% of the average parrotfish landings from 1999 to 2005. Commercial ACLs for Puerto Rico and St. Thomas were set at 52,737 and 42,500 lbs, respectively. ACLs were set below the acceptable biological catch set by the Statistical and Scientific Committee to account for uncertainty, ecological factors, and other concerns. The proposed ACLs for Puerto Rico and St. Thomas are similar to the average landings from 2006 to 2008; however, St. Croix's ACL (240,000 lbs) is lower than the average landings from 2006 to 2008 (402,744 lbs). Thus, modifications to harvest strategies are needed to reduce landings to ensure the ACL is not exceeded. This report evaluates reductions in commercial landings projected from imposing minimum and maximum size limits, slot limits, and trip limits.

Data Sources

Two different datasets were used to estimate landings reductions associated with a variety of management schemes: 1) the National Marine Fisheries Service Southeast Fisheries Science Center's (SEFSC) Trip Interview Program (TIP) and 2) the commercial landings data for the U.S. Caribbean, which consists of the USVI Department of Planning and Natural Resources

(DPNR) commercial catch record (CCR) and Puerto Rico's Department of Natural and Environmental Resources (DNR) sales receipts. TIP data were accessed on January 5, 2011 from the SEFSC's data files. These data were collected by port samplers that interviewed fishermen and measured their catch. TIP provides information on the length and numbers of parrotfish species landed, gear used, and information on the fishing trip (e.g., date sampled, location sampled, and a variety of information on fishing effort (e.g. soak time, number of lines, number of traps). In some cases the TIP sampling may only provide a sub-sample of the total catch. USVI CCR and PR DNR trip data were accessed on March 17, 2011, from the SEFSC's data files, and provided fishermen reported catch for each trip, and included landings (in pounds whole weight) by general family name (parrotfish) with information on the fishing trip (i.e., date landed or sold, location fished, and capture gear). Landings data by individual parrotfish species are not available in the USVI or Puerto Rico, but have been recommended as part of the U.S. Caribbean Commercial Data Improvement project recently undertaken by the SEFSC, USVI DPNR, Puerto Rico DNR, NOAA Fisheries Service, CFMC and territory and commonwealth fishers.

Methods

Size Limits

Size limit analyses were conducted using available U.S. Caribbean TIP data for each island (St. Croix and St. Thomas = 2008-2010; Puerto Rico = 2009-2011). Samples were aggregated across years by island to increase sample sizes when determining reductions in landings associated with various size limits, and details on pooling the data across years are provided in **Appendix A**. TIP length data were recorded using fork length (*FL*) or total length (*TL*) in cm. Since the majority of the parrotfish data were in fork length (91%), all total lengths were converted to fork length using equation 1:

$$FL = \alpha * TL \quad (1)$$

where *FL* is fork length in cm, α is a conversion parameter, and *TL* is total length in cm. Although landings data have not always been reported to species level for parrotfish, routinely the TIP data provide information on species. Data from TIP indicates seven different parrotfish species have been recorded in the samples. Length conversion parameters are shown in **Table 1**. Princess (*Scarus taeniopterus*), striped (*Scarus croicensis*) and redband (*Sparisoma aurofrenatum*) parrotfish did not require a conversion because they have truncated tails (i.e. fork length equals total length).

Table 1.- Total length to fork length conversion parameters for four parrotfish species using the equation $FL = \alpha * TL$.

Common Name	<i>Genus species</i>	α	Source
Queen Parrotfish	<i>Scarus vetula</i>	0.967	Fishbase on 4/22/2011
Redfin Parrotfish	<i>Sparisoma rubripinne</i>	0.920	Fishbase on 4/22/2011
Redtail Parrotfish	<i>Sparisoma chrysopterus</i>	0.901	Molina (2005)
Stoplight Parrotfish	<i>Sparisoma viride</i>	0.903	Fishbase on 4/22/2011

After all lengths were standardized to fork length they were converted to weight following equation 2:

$$WT = aFL^b \quad (2)$$

where WT is weight in grams, a and b are weight-length conversion parameters. Weight-length conversion parameters are shown in **Table 2**. All weight conversions are in whole weight.

Table 2.- Fork length to total weight conversion parameters for seven parrotfish species using the equation: $WT=a(FL)^b$ where weight is grams and fork length is in mm, except for queen parrotfish where the fork length is in cm.

Common Name	Genus species	a	b	Source
Princess Parrotfish	<i>Scarus taeniopterus</i>	0.010	1.875	Bohnsack and Harper (1988)
Queen Parrotfish	<i>Scarus vetula</i>	0.025	2.921	Marks and Klomp (1988)
Redband Parrotfish	<i>Sparisoma aurofrenatum</i>	8.4E-05	2.744	Bohnsack and Harper (1988)
Redfin Parrotfish	<i>Sparisoma Rubripinne</i>	1.3E-05	3.064	Bohnsack and Harper (1988)
Redtail Parrotfish	<i>Sparisoma chrysopterus</i>	0.00089	2.319	Bohnsack and Harper (1988)
Stoplight Parrotfish	<i>Sparisoma viride</i>	3.7E-05	2.905	Bohnsack and Harper (1988)
Striped Parrotfish	<i>Scarus croicensis</i>	1.3E-05	3.055	Bohnsack and Harper (1988)

All the fork lengths were converted to inches and proportional reductions in landings were estimated for a range of minimum size limits, maximum size limits, and slot limits from 8 to 15 inches FL. The size limit analysis followed the method of Chih (2003) but did not include measurement error or release mortality. Measurement error information was not available because the CFMC has never imposed size limits on parrotfish, and no information is currently available for release mortality of parrotfish of the U.S. Caribbean. Release mortality was assumed to be zero and support for imposing this assumption seems reasonable since the majority of the parrotfish landings are caught with diving (spear) and trap gears, and gill nets are prohibited. Spears are thought to have very little bycatch of parrotfish (Soldo et al. 2007) and the traps were deployed primarily in shallow water (<30 feet).

The percent reduction of parrotfish from the size limits were calculated in terms of weight by using the corresponding weight for each fish's length. The weight of parrotfish below a minimum size limit was calculated using equation 3:

$$R_1 = C - G \quad (3)$$

where R_1 is the summed weight of all fish below a size limit, C is the total weight of all the fish measured, and G is the weight of fish measured that were greater than or equal to the minimum size limit. The weight of parrotfish above a maximum size limit was calculated using equation 4:

$$R_2 = C - H \quad (4)$$

where R_2 is the summed weight of all the fish above a size limit, C is the total weight of fish measured, and H is the weight of fish measured that were less than the maximum size limit. The

percent reductions in landings were calculated for a given minimum size limit (MN), maximum size limit (MX), and slot limit using equations 5-7:

$$MN = (R_1/C) * 100 \quad (5)$$

$$MX = (R_2/C) * 100 \quad (6)$$

$$SL = MN + MX \quad (7)$$

where MN is the percent reduction from a minimum size limit, MX is the percent reduction from a maximum size limit, SL is the percent reduction in landings from a slot limit, R_1 is the weight of parrotfish below a minimum size limit, R_2 is the weight of parrotfish above a maximum size limit, and C is the total weight of parrotfish measured in the TIP database.

The percent reductions for each island were calculated with respect to gear then weighted by the percentage of commercial landings by gear. This weighted reduction was calculated with equations 8-10:

$$WRMN = \sum_{i=1}^n MNG_i * LG_i \quad (8)$$

$$WRMX = \sum_{i=1}^n MXG_i * LG_i \quad (9)$$

$$WRSL = WRMN + WRMX \quad (10)$$

where $WRMN$ is the weighted percent reduction in landings for a minimum size limit, MNG is the percent reduction from the minimum size limits for gear i , LG is the percentage of landings caught by gear i , $WRMX$ is the weighted percent reduction in landings for a maximum size limit, MXG is the maximum size limit percent reduction for gear i , and $WRSL$ is the weighted percent reduction for a slot limit.

Trip Limits

The U.S. Caribbean's USVI CCR and PR DNR trip ticket landings data from 2000 to 2008 contained 39,815 parrotfish records in St. Croix, 18,223 parrotfish records in St. Thomas, and 22,690 parrotfish records in Puerto Rico (80,728 total). The U.S. Caribbean landings data were used to evaluate reductions in landings from trip limits. A range of trip landings limits were considered ranging from 10 to 500 pounds. The estimated reduction in catch per trip was estimated using equation 11:

$$P = T - L \quad (11)$$

where P is the projected landings removed for a given trip limit, T is the total landings for a trip, and L is the imposed trip landings limit. All three variables are in pounds whole weight. P and T were summed for all trips and percent reductions associated with trip limits (PRT) were calculated using equation 12:

$$PRT = (P/T)*100. \quad (12)$$

The trip limit percent reductions for each island were calculated with respect to gear then weighted by the percentage of commercial landings by gear. This weighted reduction was calculated with equation 13:

$$WRTL = \sum_{i=1}^n PRT_i * LG_i \quad (13)$$

where $WRTL$ is the weighted percent reduction in landings for a trip limit, PRT is the percent reduction from the trip limits for gear i , and LG is the percentage of landings caught by gear i .

Combination Size and Trip Limits

Percent reductions in landings were calculated for a combination of size/slot limits and trip limits using equations 14-16:

$$CR_1 = 1 - (1 - WRMN) * (1 - WRTL) \quad (14)$$

$$CR_2 = 1 - (1 - WRMX) * (1 - WRTL) \quad (15)$$

$$CR_3 = 1 - (1 - WRSL) * (1 - WRTL) \quad (16)$$

where CR_1 is the combined weighted percent reductions in landings from both the minimum size limit and trip limit, CR_2 is the combined weighted percent reductions in landings from both the weighted maximum size limit and trip limit, and CR_3 is the combined weighted percent reductions in landings from both the slot limit and trip limit.

Results

TIP Catch Length Sample Data

The majority of U.S. Caribbean TIP parrotfish samples from 2000-2010 were from St. Croix (n=16,683) followed by Puerto Rico (n=15,360, data was available for 2011) then St. Thomas (n=1,591). Rainbow (n=44), blue (n=10), and midnight parrotfish (n=2), were removed from the dataset since their harvest is prohibited (Amendment 5). Six redband and two redband parrotfish were removed because they had lengths greater than 10 percent larger than the maximum size reported by Robins et al. (1986), and were greater than 3 standard deviations from the mean parrotfish length for the species. Records not identified to a specific parrotfish species (n=426) were also removed.

USVI CCR and PR trip ticket landings data

The U.S. Caribbean landings dataset from 2000 to 2008 included 39,815 parrotfish records in St. Croix, 18,223 parrotfish records in St. Thomas, and 22,690 parrotfish records in Puerto Rico (80,728 total). Available trip ticket data indicates U.S. Caribbean parrotfish landings from 2000-2008 peaked in 2006 (**Figure 1**). The gears that produced the majority of the landings were diving, traps, and gill nets (**Figure 2**). Despite the prohibition of gill and trammel nets in 2005 (50 CFR Parts 600 and 622) there still were landings from these gears in 2006 and 2007.




Figure 1.- Annual parrotfish landings for each U.S. Caribbean island from 2000 to 2008 (Source U.S. Caribbean landings dataset).




Figure 2.- Annual parrotfish landings from all U.S. Caribbean islands by gear type from 2000 to 2008 (Source U.S. Caribbean landings dataset). The “Other” category is composed of cast nets, spears, and unknown gear types.

Size Limit Analysis

St. Croix

Size limit analyses for St. Croix were based on the last three years of TIP data (2008 to 2010). The data from this period produced 59% of the TIP records and 35% of the TIP samples across the time period 2000 to 2010. Also, these years are more likely to represent current harvesting practices. A large reduction in landings (>50%) was estimated for either minimum size limits ≥ 12 inches FL or maximum size limits ≤ 11 inches FL (**Table 3**). **Table 4** provides a range of estimated percent reductions in landings for several combinations of minimum and maximum size limits (i.e., ‘slot limits’).

Table 3.- Percent reductions in parrotfish landings (pounds and percentage) for minimum and maximum size limits applied to St. Croix TIP data for the years 2008 to 2010. The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “lbs” in the minimum size column represent variable G in equation 3, and the numbers listed under “lbs” in the maximum size column represent variable H for equation 4. Variable C in equations 3 and 4 is the “No Limit” value (5,496 pounds).

Size Limit (inches FL)	Minimum size		Maximum Size	
	lbs	%	lbs	%
No Limit	5,496		5,496	
8	24	0.4	5,472	99.6
9	319	5.8	5,177	94.2
10	997	18.1	4,499	81.9
11	2,371	43.1	3,125	56.9
12	4,055	73.8	1,441	26.2
13	5,022	91.4	474	8.6
14	5,329	97.0	167	3.0
15	5,425	98.7	71	1.3

Table 4.- Weighted percent reductions in parrotfish landings (in lbs) for various slot limits created from St. Croix TIP data for the years 2008 to 2010.

Minimum Size (inches)	Maximum Size (inches)							
	8	9	10	11	12	13	14	15
8	-	94.6	82.3	57.3	26.7	9.1	3.5	1.7
9		-	87.7	62.7	32.0	14.4	8.8	7.1
10			-	75.0	44.4	26.8	21.2	19.4
11				-	69.4	51.8	46.2	44.4
12					-	82.4	76.8	75.1
13						-	94.4	92.7
14							-	98.2

St. Thomas

Size limit analyses for St. Thomas were based on the last three years of TIP data (2008 to 2010). This time period had the majority of the TIP records (61%) and samples (68%) from 2000 to 2010. Also, these years likely represent current harvesting conditions. A large reduction in landings (>50%) was estimated for minimum size limits ≥ 12 inches FL or maximum size limits ≤ 11 inches FL (**Table 5**). **Table 6** provides a range of estimated percent reductions in landings for combinations of minimum and maximum size limits (i.e., ‘slot limits’).

Table 5.- Percent reductions in parrotfish landings (pounds and percentage) for minimum and maximum size limits applied to St. Thomas TIP data for the years 2008 to 2010. The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “lbs” in the minimum size column represent variable G in equation 3, and the numbers listed under “lbs” in the maximum size column represent variable H for equation 4. Variable C in equations 3 and 4 is the “No Limit” value (1,059 pounds).

Size Limit (inches FL)	Minimum size		Maximum Size	
	Lbs	%	lbs	%
No Limit	1,059		1,059	
8	2	0.1	1,057	99.9
9	17	1.6	1,042	98.4
10	136	12.8	923	87.2
11	474	44.7	585	55.3
12	837	79.0	222	21.0
13	1,008	95.2	51	4.8
14	1,041	98.3	18	1.7
15	1,051	99.2	8	0.8

Table 6.- Weighted percent reductions in parrotfish landings for various slot limits created from St. Thomas TIP data for years 2008 to 2010.

Minimum Size (inches)	Maximum Size (inches)							
	8	9	10	11	12	13	14	15
8	-	98.5	87.3	55.4	21.1	4.9	1.8	0.9
9		-	88.8	56.9	22.6	6.4	3.3	2.4
10			-	68.1	33.8	17.6	14.5	13.6
11				-	65.7	49.5	46.4	45.5
12					-	83.8	80.7	79.8
13						-	96.9	96.0
14							-	99.1

Puerto Rico

Size limit analyses were conducted using 2009-2011 TIP data, which accounted for 15% of the total TIP records and 16% of the total parrotfish records during 2000-2011. Large reduction in landings (>50%) were estimated for either minimum size limits ≥ 12 inches FL or maximum size limits ≤ 11 inches FL (**Table 7**). **Table 8** provides a range of estimated percent reductions in landings for various combinations of minimum and maximum size limits (i.e., ‘slot limits’).

Table 7.- Percent reductions in parrotfish landings (pounds and percentage) for minimum and maximum size limits applied to Puerto Rico TIP data for the years 2009 to 2011. The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “lbs” in the minimum size column represent variable G in equation 3, and the numbers listed under “lbs” in the maximum size column represent variable H for equation 4. Variable C in equations 3 and 4 is the “No Limit” value (2,377 pounds).

Size Limit (inches FL)	Minimum size		Maximum Size	
	lbs	%	lbs	%
No Limit	2,377		2,377	
8	21	0.9	2,356	99.1
9	154	6.5	2,223	93.5
10	531	22.3	1,846	77.7
11	1072	45.1	1,305	54.9
12	1692	71.2	685	28.8
13	2178	91.6	199	8.4
14	2257	94.9	120	5.1
15	2270	95.5	107	4.5

Table 8.- Weighted percent reductions in parrotfish landings for various slot limits created from Puerto Rico TIP data for years 2009 to 2011.

Minimum Size (inches)	Maximum Size (inches)							
	8	9	10	11	12	13	14	15
8	-	94.4	78.6	55.8	29.7	9.2	5.9	5.4
9		-	84.2	61.4	35.3	14.8	11.5	11.0
10			-	77.2	51.2	30.7	27.4	26.8
11				-	73.9	53.5	50.2	49.6
12					-	79.5	76.2	75.7
13						-	96.7	96.1
14							-	99.4

Size limit restrictions have the potential to impact each parrotfish species in a different way. Relative impacts depend primarily on the size distribution and population abundance of each parrotfish species. The most common parrotfish species in the most recent available TIP data for all three areas (St. Croix 2008-2010, St. Thomas 2008-2010, Puerto Rico 2009-2011) were redband, stoplight, and redband parrotfish (**Table 9**). A box plot of the size composition for each species (**Figure 3**) shows the majority of the parrotfish lengths in the catch are relatively similar.

Table 9.- Percentage of TIP records by species for each island for the most recent data (St. Croix 2008-2010, St. Thomas 2008-2010, and Puerto Rico 2009-2011). Numbers in parentheses are sample size.

	Princess	Queen	Redband	Redfin	Redtail	Stoplight	Striped
St. Croix	1.4 (87)	0.2 (13)	7.9 (463)	5.6 (329)	49.4 (2,907)	35.5 (2,090)	0.0 (0)
St. Thomas	0.7 (7)	0.2 (2)	0.8 (8)	3.6 (38)	45.8 (485)	48.9 (518)	0.1 (1)
Puerto Rico	3.0 (225)	1.3 (100)	3.7 (281)	3.6 (275)	47.6 (3,605)	40.7 (3,079)	0.02 (2)
All Three Islands	2.2 (319)	0.8 (115)	5.2 (752)	4.4 (642)	48.2 (6,997)	39.2 (5,687)	0.02 (3)




Figure 3.- Box plot of the size of each parrotfish species created from the most recent TIP data for all three islands (St. Croix 2008-2010, St. Thomas 2008-2010, and Puerto Rico 2009-2011).

Striped parrotfish were not included in the boxplot because there was only one record in the most recent data.

Trip Limit Analysis

St. Croix

The annual landings data from St. Croix (2007 and 2008) accounted for 19% of the records and 21% of the landings by weight (lbs) between 1998 and 2008. The average, minimum, and maximum parrotfish landings per trip were 86, 1, and 2,996 pounds for 2007, and 78, 1, and 905 pounds for 2008. A trip limit analysis estimated large reductions in landings (>50%) occurred at trip limits of 60 pounds or less (**Table 10**). Trip limits imposed in St. Croix would have the greatest impact on parrotfish collected with dive gear (**Table 11**), which may also include spear and net gears.

Table 10.- Percent reductions in parrotfish landings (pounds and percentage) for various trip limits applied to St. Croix landings data for 2007 and 2008 (n=9,449 trips). The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “Landings (lbs) Reduction” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value of 774,888 pounds.

Trip Limit	Landings (lbs) Reduction	Percent Reduction
No Limit	774,888	
10	683,700	88.2
20	604,992	78.1
30	539,216	69.6
40	482,008	62.2
50	432,091	55.8
60	389,124	50.2
70	352,723	45.5
80	321,209	41.5
90	293,796	37.9
100	269,509	34.8
150	194,548	25.1
200	140,732	18.2
300	59,437	7.7
400	16,866	2.2
500	5,533	0.7

Table 11.- Percent reductions in parrotfish landings (pounds and percentage) for each gear based on St. Croix landings data for 2007 and 2008. The numbers listed under “Landed (lbs)” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value for each gear.

Trip Limits (lbs)	Castnet		Diving		Traps		Line Fishing		Seine Net		Trammel Net		Gillnet	
	Landed (lbs)	%												
	Reduction		Reduction		Reduction		Reduction		Reduction		Reduction		Reduction	
No Limit	1,906		491,820		104,589		22,333		55,559		45,128		53,326	
10	1,466	76.9	432,322	87.9	83,034	79.4	17,705	79.3	54,089	97.4	43,858	97.2	50,971	95.6
20	1,034	54.2	376,252	76.5	69,704	66.6	14,030	62.8	52,619	94.7	42,588	94.4	48,654	91.2
30	630	33.1	327,423	66.6	61,115	58.4	11,083	49.6	51,149	92.1	41,318	91.6	46,364	86.9
40	312	16.4	284,411	57.8	54,969	52.6	8,322	37.3	49,689	89.4	40,048	88.7	44,084	82.7
50	131	6.9	247,250	50.3	49,877	47.7	5,894	26.4	48,229	86.8	38,778	85.9	41,813	78.4
60	48	2.5	216,149	43.9	45,058	43.1	3,877	17.4	46,769	84.2	37,508	83.1	39,556	74.2
70	5	0.3	190,690	38.8	40,446	38.7	2,555	11.4	45,309	81.6	36,238	80.3	37,306	70.0
80	0	0.0	169,473	34.5	36,005	34.4	1,762	7.9	43,849	78.9	34,968	77.5	35,066	65.8
90	0	0.0	151,615	30.8	31,814	30.4	1,292	5.8	42,389	76.3	33,698	74.7	32,876	61.7
100	0	0.0	136,225	27.7	28,146	26.9	963	4.3	40,939	73.7	32,428	71.9	30,696	57.6
150	0	0.0	95,471	19.4	18,069	17.3	345	1.5	33,789	60.8	26,178	58	20,646	38.7
200	0	0.0	70,056	14.2	11,305	10.8	80	0.4	26,758	48.2	19,988	44.3	12,471	23.4
300	0	0.0	30,397	6.2	4,095	3.9	0	0.0	13,959	25.1	9,041	20.0	1,991	3.7
400	0	0.0	6,277	1.3	1,550	1.5	0	0.0	6,395	11.5	2,553	5.7	70	0.1
500	0	0.0	2,496	0.5	800	0.8	0	0.0	3,620	6.5	1,069	2.4	0	0.0

St. Thomas

The annual landings data from St. Thomas (2007 and 2008) accounted for 18% of the records and landings in weight (lbs) during 2000-2008. The average, minimum, and maximum parrotfish landings per trip were 24, 1, and 145 pounds for 2007, and 24, 2, and 200 pounds for 2008. A trip limit analysis estimated large reductions in landings (>50%) occurred at a trip limit of 10 pounds (**Table 12**). Trip limits imposed in St. Thomas would have the greatest impact on parrotfish collected with traps (**Table 13**).

Table 12.- Percent reductions in parrotfish landings (pounds and percentage) for various trip limits applied to St. Thomas landings data for 2007 and 2008 (n=3,366 trips). The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “Landings (lbs) Reduction” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value of 79,788 pounds.

Trip Limit	Landings (lbs) Reduction	Percent Reduction
No Limit	79,788	
10	50,073	62.8
20	33,291	41.8
30	22,794	28.5
40	15,345	19.2
50	9,726	12.2
60	5,564	7.0
70	3,128	3.9
80	1,871	2.3
90	1,247	1.6
100	935	1.2
150	156	0.2
200	0	0.0
300	0	0.0
400	0	0.0
500	0	0.0

Table 13.- Percent reductions in parrotfish landings (pounds and percentage) for each gear based on St. Thomas landings data for 2007 and 2008. The numbers listed under “Landed (lbs)” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value for each gear.

Trip Limits (lbs)	Diving		Traps		Line Fishing		Seine Net	
	Landed (lbs)	%						
	Reduction		Reduction		Reduction		Reduction	
No Limit	827		77,944		800		213	
10	491	59.4	49,212	63.1	259	32.4	137	64.3
20	252	30.5	32,850	42.1	142	17.8	80	37.6
30	132	16.0	22,494	28.9	85	10.6	50	23.5
40	65	7.9	15,161	19.5	60	7.5	20	9.4
50	20	2.4	9,641	12.4	40	5.0	0	0.0
60	10	1.2	5,531	7.1	20	2.5	0	0.0
70	0	0.0	3,083	4.0	10	1.3	0	0.0
80	0	0.0	1,863	2.4	0	0.0	0	0.0
90	0	0.0	1,283	1.6	0	0.0	0	0.0
100	0	0.0	930	1.2	0	0.0	0	0.0
150	0	0.0	165	0.2	0	0.0	0	0.0
200	0	0.0	0	0.0	0	0.0	0	0.0
300	0	0.0	0	0.0	0	0.0	0	0.0
400	0	0.0	0	0.0	0	0.0	0	0.0
500	0	0.0	0	0.0	0	0.0	0	0.0

Puerto Rico

The annual landings data (2008 and 2009) from Puerto Rico accounted for 13% of the records and 8% of the landings by weight (lbs) during 1998 to 2009. The average, minimum, and maximum parrotfish landings per trip were 51, 2, and 5,000 pounds for 2008, and 31, 1, and 1,383 pounds for 2009. A trip limit analysis estimated large reductions in landings (>50%) occurred at a trip limit of 10 pounds (**Table 14**). Trip limits imposed in Puerto Rico would have the greatest impact on parrotfish collected with traps (**Table 15**).

Table 14.- Percent reductions in parrotfish landings (pounds and percentage) for various trip limits applied to Puerto Rico landings data for 2008 and 2009 (n=3,709 trips). The reductions were calculated in terms of parrotfish weight (lbs) with respect to gear. Then the reductions were weighted by the percentage of landings for each gear type. The numbers listed under “Landings (lbs) Reduction” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value of 56,487 pounds.

Trip Limit lbs	Landings (lbs) Reduction	Percent Reduction
No Limit	56,487	
10	29,309	54.6
20	18,104	35.2
30	12,899	25.6
40	9,515	19.2
50	7,112	14.6
60	5,373	11.2
70	4,092	8.6
80	3,146	6.7
90	2,469	5.3
100	1,961	4.3
150	716	1.8
200	385	1.0
300	0	0
400	0	0
500	0	0

Table 15.- Percent reductions in parrotfish landings (pounds and percentage) for each gear based on Puerto Rico landings data for 2008 and 2009. The numbers listed under “Landed (lbs)” represent variable P in equation 11. Variable T in equation 11 is the “No Limit” value for each gear.

Trip Limits (lbs)	Diving		Traps		Line Fishing		Seine Net		Trammel Net		Gillnet	
	Landed (lbs)	% Reduction										
No Limit	16,148		13,238		13,253		1,669		6,833		5,346	
10	7,668	47.5	4,868	36.8	8,463	63.9	895	53.6	5,172	75.7	2,243	42.0
20	4,101	25.4	2,032	15.3	6,100	46.0	488	29.2	3,883	56.8	1,500	28.1
30	2,732	16.9	1,088	8.2	4,637	35.0	369	22.1	2,953	43.2	1,120	21.0
40	1,987	12.3	485	3.7	3,630	27.4	309	18.5	2,224	32.5	880	16.5
50	1,460	9.0	148	1.1	2,882	21.7	250	15.0	1,672	24.5	700	13.1
60	1,049	6.5	26	0.2	2,274	17.2	200	12.0	1,254	18.4	570	10.7
70	776	4.8	5	0.0	1,776	13.4	150	9.0	942	13.8	443	8.3
80	575	3.6	0	0.0	1,437	10.8	105	6.3	691	10.1	338	6.3
90	419	2.6	0	0.0	1,192	9.0	70	4.2	531	7.8	257	4.8
100	299	1.9	0	0.0	1,014	7.7	50	3.0	397	5.8	201	3.8
150	21	0.1	0	0.0	585	4.4	0	0.0	100	1.5	10	0.2
200	0	0.0	0	0.0	335	2.5	0	0.0	50	0.7	0	0.0
300	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
400	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
500	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

Combination Size and Trip Limit Analyses

Minimum size, maximum size, and slot limit results were combined with trip limit results to determine percent reductions in landings for each island (Tables 16, 17, 18, and 19). A range of sizes ranging from 8 to 14 inches FL were chosen to provide a variety of estimated percent reductions in landings. Trip limits of 10 to 50, 100, 150, and 200 pounds were chosen because they provided a variety of estimated percent reductions in landings for each island (Tables 10, 12, and 14).

Table 16.- Weighted percent reductions in parrotfish landings for minimum and maximum size limits combined with various trip limits for St. Croix.

		Trip Limit (lbs)							
		10	20	30	40	50	100	150	200
	8	88.3	78.2	69.8	62.5	56.1	35.2	25.6	18.7
	9	89.0	79.6	71.7	64.8	58.9	39.3	30.3	23.9
Min Size (in)	10	90.5	82.4	75.5	69.6	64.4	47.5	39.7	34.1
	11	93.2	87.5	82.6	78.3	74.7	62.6	57.1	53.1
	12	97.0	94.4	92.2	90.3	88.6	83.2	80.7	78.9
	10	97.7	95.7	94.1	92.6	91.4	87.3	85.4	84.1
	11	95.0	90.6	87.0	83.9	81.1	72.2	68.0	65.1
Max Size (in)	12	91.2	83.7	77.4	71.9	67.2	51.6	44.4	39.3
	13	89.1	79.8	71.9	65.1	59.2	39.8	30.8	24.4
	14	88.5	78.7	70.4	63.2	57.0	36.5	27.1	20.3

Table 17.- Weighted percent reductions in parrotfish landings for minimum and maximum size limits combined with various trip limits for St. Thomas.

		Trip Limit (lbs)							
		10	20	30	40	50	100	150	200
	8	62.9	41.9	28.6	19.3	12.3	1.3	0.3	0.1
	9	63.4	42.7	29.7	20.5	13.6	2.8	1.8	1.6
Min Size (in)	10	67.6	49.3	37.7	29.6	23.5	13.9	13.0	12.8
	11	79.4	67.8	60.5	55.3	51.5	45.4	44.8	44.7
	12	92.2	87.8	85.0	83.1	81.6	79.3	79.1	79.0
	10	95.2	92.5	90.8	89.6	88.7	87.3	87.2	87.2
	11	83.4	74.0	68.0	63.9	60.7	55.8	55.4	55.3
Max Size (in)	12	70.6	54.0	43.5	36.1	30.6	21.9	21.1	21.0
	13	64.6	44.6	31.9	23.1	16.4	5.9	5.0	4.8
	14	63.4	42.8	29.7	20.6	13.7	2.9	1.9	1.7

Table 18.- Weighted percent reductions in parrotfish landings for minimum and maximum size limits combined with various trip limits for Puerto Rico.

		Trip Limit (lbs)							
		10	20	30	40	50	100	150	200
	8	54.8	35.6	26.0	19.7	15.1	4.9	2.4	1.6
	9	56.0	37.2	27.8	21.7	17.2	7.3	4.8	4.0
Min Size (in)	10	60.6	43.8	35.5	30.0	26.0	17.0	14.9	14.2
	11	68.5	55.1	48.4	44.0	40.8	33.7	31.9	31.4
	12	81.4	73.4	69.5	66.9	65.0	60.8	59.8	59.4
	10	94.0	91.4	90.1	89.3	88.7	87.3	87.0	86.8
	11	86.1	80.1	77.2	75.2	73.8	70.7	69.9	69.6
Max Size (in)	12	73.2	61.8	56.1	52.3	49.6	43.5	42.1	41.6
	13	60.8	44.0	35.7	30.3	26.3	17.4	15.2	14.5
	14	56.3	37.6	28.4	22.3	17.9	7.9	5.5	4.7

Table 19.- Weighted percent reductions in parrotfish landings for slot limits and trip limits for all three islands.

	Slot Limit (inches)	Trip Limit (lbs)							
		10	20	30	40	50	100	150	200
St. Croix	8 to 12	91.3	83.9	77.6	72.2	67.5	52.0	44.9	39.8
	8 to 14	88.6	78.8	70.6	63.4	57.3	36.9	27.6	20.9
	10 to 14	90.8	82.9	76.3	70.5	65.6	49.2	41.6	36.2
St. Thomas	8 to 12	70.7	54.1	43.6	36.3	30.7	22.1	21.3	21.1
	8 to 14	63.5	42.9	29.8	20.7	13.8	3.0	2.1	1.9
	10 to 14	68.2	50.3	38.9	31.0	25.0	15.6	14.7	14.6
Puerto Rico	8 to 12	73.5	62.1	56.5	52.8	50.1	44.1	42.6	42.1
	8 to 14	56.6	38.0	28.8	22.7	18.3	8.5	6.1	5.3
	10 to 14	62.3	46.2	38.3	33.0	29.2	20.6	18.6	17.9

Discussion

The CFMC recently implemented ACLs for all federally managed species. Based on ACLs recommended for parrotfish, reductions in landings will likely be required in St. Croix to prevent the ACL from being exceeded, and accountability measures from being triggered. The ACL for St. Croix is 40% lower than the average landings for 2006-2008 (CFMC 2010). Assuming previous year's landings are a reasonable predictor of future landings trends, a 40% reduction in landings could be accomplished with a minimum or maximum size of 11 inches (**Table 3**). These management measures could aid in decreasing the landings below the recommended ACL level given all other factors remaining constant (e.g., fisher behavior, fishing effort and distribution).

Evaluation of size limit data in these analyses suggested most parrotfish sampled through the TIP program were ≤ 12 inches FL (**Figure 3**). Size limits can potentially reduce harvest as well as reduce fishing mortality, protect immature fish and increase yield-per-recruit from a fish stock, assuming discard mortality is low. Size limits can also minimize recruitment overfishing, which can lead to a stock biomass level below maximum yield. Minimum size limits have the potential to increase productivity of a stock by allowing the fish to spawn before being harvested. Parrotfish become mature at a range of lengths varying spatially and across species. A literature review provided a range of lengths at maturity for U.S. Caribbean parrotfish from ~6 to 11 inches FL (Robertson and Warner 1978; Koltes 1993). A minimum size limit has the potential to improve the redband parrotfish stock because the majority of their harvests were below the size at maturity (~9.25 inches, Randall 1963) (**Figure 3**).

Another consideration of minimum size limits is the potential to cause fishermen to remove the larger fish, which produce exponentially more eggs than smaller fish (Bohnsack, 1990). Also, the absence of larger and older spawning stock may lower recruitment by preventing parrotfish from living long enough to survive through periods when conditions are poor for offspring survival (Hawkins and Roberts 2003), especially since the maximum ages of parrotfish species in the current catch range from only 3 to 7 years (Molina-Urena 2005). An additional consideration is all Caribbean parrotfish are protogynous hermaphrodites, changing sex from females to males. Generally, minimum size limits will protect the females but shift fishing pressure upon the males. This makes them highly susceptible to overfishing. Chronic removal of the male

parrotfish will dramatically decrease sperm availability and act as a limiting factor during mating (Clua and Legendre 2008). Another consideration is the reproductive behavior since male parrotfish maintain harems. The reproduction of the entire harem will be interrupted when a male is harvested, thus seriously reducing the reproductive output as seen by hogfish (*Lachnolaimus maximus*) (McBride and Johnson 2007). Also, parrotfish must reach a genetically determined size threshold before they can become males (Warner 1988; Clua and Legendre 2008). If fishing prevents individuals from growing large enough to undergo sex change then the population could become non-reproductive. Hawkins and Roberts (2003) examined parrotfish populations in Jamaica and the Dominican Republic and discovered fishing pressure eliminated stoplight and queen parrotfish males to the point where the population depends on recruitment from distant sources. Minimum size limits also have the potential to alter the balance of the species since each parrotfish species will be impacted differently. This could be counterproductive to ecosystem health.

Implementing a maximum size limit could help prevent the removal of older, larger fish in the population. Larger parrotfish are more likely to be male. Imposing a maximum size limit increases male survival and could prevent interruption of spawning harems. A maximum size limit of 11 inches for St. Croix was estimated to reduce landings by 42% (Table 1) and can, potentially, decrease the current landings below the proposed ACL level. However, the continued removal of small fish, which are primarily females, could hamper spawning success. Additionally, a maximum size limit will only lead to a sustainable fishery if individuals live long enough to reach the maximum size limit.

Any minimum and maximum size limits need to consider socioeconomic impacts. There may be a range of parrotfish sizes that are more preferred in the markets.

There are a range of slot limits that can potentially reduce St. Croix parrotfish landings to the ACL level, which are highlighted in Table 20. In addition to the potential impacts of minimum and maximum size limits discussed earlier, a slot limit has the potential to protect females with a minimum size limit and the males with a maximum size limit.

Table 20.- Weighted percent reduction of parrotfish landings for various slot limits created from St. Croix TIP data for years 2008 to 2010. Cells highlighted in yellow have a reduction in landings of 40 percent or greater.

Minimum Size (inches)	Maximum Size (inches)							
	8	9	10	11	12	13	14	15
8	-	94.6	82.3	57.3	26.7	9.1	3.5	1.7
9		-	87.7	62.7	32.0	14.4	8.8	7.1
10			-	75.0	44.4	26.8	21.2	19.4
11				-	69.4	51.8	46.2	44.4
12					-	82.4	76.8	75.1
13						-	94.4	92.7
14							-	98.2

Parrotfish have become one of the most important reef fish species-groups harvested in St. Croix (Toller 2007). The use of gill and trammel nets was prohibited in St. Croix in 2006 in both state and federal waters, however these gears were used after 2006 (**Figure 2**). Additionally, 70% of scuba landings are mis-reported (Toller 2007) because gill nets were used in combination with scuba gear. Toller (2007) analyzed St. Croix parrotfish landings and determined a cut-off threshold of 162.5 pounds to separate net-scuba from scuba-only landings. Therefore, a potential regulation to eliminate illegal netting is to set a trip limit below 162 pounds. A 150 pound trip limit can potentially reduce landings by 25%.

Finally, the results are based on the data available and assumptions used. Analyses assume the parrotfish length frequency distribution of the TIP dataset is representative of the catch, however, the TIP dataset has some caveats. The sampling depends on volunteer fishermen allowing a biologist to measure and count their catch. This volunteer option may bias the TIP data and could prevent the dataset from being an accurate representation of the overall catch. Additionally, fishermen involved in illegal activity will likely not participate. This issue was discussed at a SEDAR workshop in January of 2009 addressing non-reporting as a significant concern for Caribbean catch statistics. Another limitation of the TIP data is flaws in the sampling design with nonrandom sampling potentially biasing the results.

Analyses also assume fishermen will fully comply with size limits and no measurement error will occur. Retention of parrotfish below or above specified size limits would reduce the reductions estimated in this report. Additionally, trip limit reductions do not account for shifts in fishing effort or other behavior changes. If fishermen take more fishing trips to compensate for more restrictive trip limits, then reductions summarized herein are overestimated. The analysis is also affected by potential biases associated with sampling, such as spatial and temporal differences in sampling intensity over time.

In summary size limits and trip limits could potentially reduce parrotfish landings to levels equal to or less than the ACL set by the CFMC. If size limits are to be considered as a measure for reducing landings to meet the ACL, then parrotfish reproductive biology should be considered as size and slot limits may differentially affect female and male parrotfish. Trip limits could reduce landings, but would decrease the efficiency of fishermen that frequently produce large landings.

References

- Bohnsack, J.A. and D.E. Harper. 1988. Length-weight relationship of selected marine reef fishes from the southeastern United States and the Caribbean. NOAA Technical Memorandum NMFS-SEFSC-215, 31 p.
- Bohnsack, J.A. 1990. The potential of marine fishery reserves for reef fish management in the US southern Atlantic. NOAA Tech. Memo NMFS-SEFSC-261, Miami, Florida.
- Brock, R.E. 1979. An experimental study on the effects of grazing by parrotfishes and role of refuges in benthic community structure. *Mar. Biol.* 51: 381-388.
- Burkepile, D.E. and M.E. Hay. Impact of herbivore identity on algal successions and coral growth on a Caribbean Reef. *Plos ONE*. 5(1): e8963. Doi:10.1371/journal.pone.0008963.
- Caribbean Fisheries Management Council. 2010. Public hearing draft amendment 2 to the fishery management plan for the queen conch fishery of Puerto Rico and the U.S. Virgin Islands and

- amendment 5 to the reef fish fishery management plan of Puerto Rico and the U.S. Virgin Islands. 468 pages.
- Chih, C.P. 2003. An evaluation of minimum size limits for managing the vermilion snapper commercial fishery. Sustainable Fisheries Division Contribution No.SFD-2003-034.
- Clua, E. and P.Legendre. 2008. Shifting dominance among Scarid species on reefs representing a gradient of fishing pressure. *Aquatic Living Resources*. 21:339-348.
- Figuerola, M., Matos-Caraballo, D., Torres, W. 1998. Maturation and reproductive seasonality of four reef fish species in Puerto Rico. In: Creswell, R.L. (ed.). Proceedings of the 50th annual session. Gulf and Caribbean Fisheries Institute, Merida, Mexico, November 1997. pp. 938-968.
- Hawkins, J.P. and C.M. Roberts. 2003. Effects on sex-changing Caribbean parrotfish. *Biological conservation*. 115:213-226.
- Koltes, K.H. 1993. Aspects of the reproductive biology and social structure of the stoplight parrotfish *Sparisoma viride*, at Grand Turk, Turks and Caicos Islands, B.W.I. *Bulletin of Marine Science*. 52(2):792-805.
- Marks, K.W. and K.D. Klomp. 2003. Appendix Two: Fish biomass equations. Pp. 625-628 in J.C. Lang (ed.), *Status of Coral Reefs in the Western Atlantic: Results of Initial Surveys, Atlantic and Gulf Rapid Reef Assessment (AGRRA) Program*. Smithsonian Atoll Research Bulletin 496.
- McBride, R.S. and M.R. Johnson. 2007. Sexual development and reproductive seasonality of hogfish (Labridae:*Lachnolaimus maximus*), an hermaphroditic reef fish. *Journal of Fish Biology*. 71:1270-1292.
- Molina-Urena, H. 2005. Towards an ecosystem approach for non-target reef fishes: habitat uses and population dynamics of South Florida parrotfishes (Perciformes: Scaridae). University of Miami, Coral Gables, Florida. Dissertation. 322 pages.
- Mumby, P.J. 2006. The impact of exploited grazers (Scaridae) on the dynamics of Caribbean Coral Reefs. *Ecological Applications*. 16(2):747-769.
- Randall, J.E. 1963. An analysis of the fish populations of artificial and natural reefs in the virgin islands. *Caribbean Journal of Science* 3(1):31-47.
- Robertson, D.R., and R.R. Warner. 1978. Sexual patterns in the Labroid fishes of the Western Caribbean. II The parrot fishes (Scaridae). *Smithsonian Contribution Zoology*. 255: 1-26.
- Robins, C.R., and G.C. Ray. 1986. *Atlantic Coast Fishes*. Peterson Field Guide. Houghton Mifflin. Boston, MA. 354 p.
- Sokal, R.R. and F.J. Rohlf. 1995. *Biometry: the principles and practice of statistics in biological research*, 3rd edition. New York: W.H. Freeman and Company. 887 pages.
- Soldo, A., F.A. Keljo, and J. Brcic. 2007. Spear fishing in the eastern Adriatic. *Rapp. Comm. Int. Mer. Medit.* 38:603.
- Soldo, A., F.A. Keljo, and J. Brcic. 2007. Spear fishing in the eastern Adriatic. *Rapp. Comm. Int. Mer. Medit.* 38:603.
- Toller, W. 2007. Revised estimate of recent parrotfish harvests by St. Croix commercial divers: separation of reported net-scuba landings from scuba-only landings. Division of Fish and Wildlife. Government of the U.S. Virgin Islands.
- Warner, R.R. 1988. Sex change in fishes: hypothesis, evidence, and objections. *Environmental Biology of Fishes*. 22:81-90.