

**STOCK ASSESSMENT AND FISHERY  
EVALUATION REPORT**

**CARIBBEAN SPINY LOBSTER FISHERY**

**1992**

**National Marine Fisheries Service  
Southeast Regional Office  
9450 Koger Boulevard  
St. Petersburg, FL 33702**

## INTRODUCTION

The Caribbean Fishery Management Council's (CFMC) Fishery Management Plan (FMP) for spiny lobster in the U.S. Caribbean was implemented on January 1, 1985. It identified a number of activities that require the attention of the National Marine Fisheries Service (NMFS) and the Caribbean Fishery Management Council (CFMC), in cooperation with the Commonwealth of Puerto Rico, and the Territory of the U.S. Virgin Islands through their pertinent agencies: Department of Natural Resources (DNR) and the Fisheries Research Laboratory in Puerto Rico, and the Department of Planning And Natural Resources in the U.S. Virgin Islands. A central management measure for this FMP is a 3.5 inch (89 mm) carapace length as the minimum legal size limit. A spiny lobster stock assessment workshop was conducted at the CFMC offices in San Juan, Puerto Rico on September 11-13, 1990 to meet FMP requirements for continual monitoring and subsequent action as data becomes available. This report is the resulting Stock Assessment And Fishery Evaluation (SAFE) Report for the spiny lobster resource in the U.S. Caribbean.

## METHODS

In preparation for the assessment, data sheets from approximately 950 trip interviews from St. Thomas, St John, and Puerto Rico from 1985 through 1989 were assembled by the CFMC staff and submitted to Miami Laboratory NMFS for data entry in the Trip Interview Program (TIP) format. Additional data sheets for three

years of sampling from 1987 through 1989 for St. Croix were entered by CFMC staff. Data sets, representing over 25,000 measured lobster, were combined for length-frequency analysis using SAS<sup>1</sup> software at the workshop. Participants examined data and conducted analyses where appropriate. The assessment team chose to use Puerto Rico, St. Croix, and the combined St. John and St. Thomas areas as appropriate units for analysis. St. Croix was separated from the other Virgin Islands because it is located on a separate geological platform.

## RESULTS AND DISCUSSION

### Data Collection, Entry, and Management

#### Available Data

Several problems were noted in data collection procedures, or in data base management, which limited the types of analyses that were possible. These problems are detailed to improve future efforts and to give other researchers examples of situations to avoid:

1. Sampling units and types were not recorded on many data sheets so confusion existed as to whether measurements were kilos or pounds, centimeters or inches, carapace length or total length, lobster traps or fish traps, etc.

---

<sup>1</sup> SAS is a registered trademark of the SAS Institute, Inc., Box 8000, Cary, North Carolina 27511-8000. The National Marine Fisheries Service and other organizations listed in this report do not endorse any particular commercial product.

2. Zero catches (i.e. trips that targeted lobster but with zero landings) were not recorded in the trip interview samples for Puerto Rico.

3. Numerous coding problems existed in the data base because data sheets and codes were not standardized between islands or between time periods within islands. Some area codes were either erroneous or were not documented. The uncertainty in how to interpret the data sheets created confusion for data entry personnel in Miami, who were not familiar with the peculiarities of the data collection program, such as sampling methodologies, exact landing locations, species codes, etc.

4. Completely and partially sampled trips were not distinguished on data sheets for Puerto Rico, which made calculating catch-per-unit-effort impossible.

5. In some cases units were recorded to several decimal places implying false precision. Apparently some measurements were collected in kilograms but converted by calculator to pounds before entry on a data sheet.

6. Virgin Island carapace measurements were recorded to the nearest tenth of an inch while in Puerto Rico measurements were to the nearest mm.

## Recommendations

As part of a solution in addressing these problems, the workshop recommends standardization of data collection and data base management. Some problems in the analysis of this data were caused by lack of standardization as to how data were collected or recorded. For example, the sample sizes and coverage of length-frequency samples from the Virgin Islands were of limited use because the measurement units (0.1 in) were too large. We recommend using 1 mm increments. In Puerto Rico only weights were frequently recorded which were less useful than if combined with length measurements. We recommend preference be given to length measurements with subsamples being weighed where possible, however, weights without lengths are preferable to no data.

Where possible, data entry should be done by the data collecting agency to avoid misinterpretation. Many of the problems encountered in interpreting data sheets could have been solved by having the organizations or individuals that collected information enter data, preferably as soon as possible after collecting information. Although all data now being collected in Puerto Rico is now being entered very soon after it is collected, much of the data used in this workshop were entered into a computer several years after collection. Most entry errors could be corrected by inspection of print-outs of records immediately after having been entered. Many errors could be corrected by error checking programs that identify unusual values.

A standardized storage format should allow basic data analysis for local governmental use as well as for more complex analysis. The recently renovated microcomputer TIP program, TIP Data Entry System Version 3.0, developed by the Southeast Fisheries Center is one possible standardization solution. This system must be successfully tested in the field and allow easy data retrieval for local uses. A Spanish language version for Puerto Rico would be most welcome.

### Fishery Trends

#### Total Landings

Total spiny lobster landings data were assembled for Puerto Rico and the Virgin Islands (Table 1). In Puerto Rico total annual landings averaged 317,451 lbs for 23 years of available data, but have fluctuated over time (Fig. 1). Total annual landings increased from 1972 to a high of 512,000 lbs in 1979, and declined from 1979 to a low of 143,761 lbs in 1988. Thus, 1988 and 1989 total landings were, respectively, only 28% and 36% of the maximum reported landings in 1979. Despite uncertainty about the accuracy of calculated values for some years (see Matos and Sadovy, 1990), the review team considered that the data probably reflected general landings trends.

Total landings averaged 36,534 lbs for St. Thomas and St. John and 7,284 for St. Croix between 1980 and 1988 (Fig. 2). Landings in the Virgin Islands appeared relatively stable during the time

that landings data were available between 1980 and 1988. Total annual landings were higher from St. Thomas/St. John than from St. Croix presumably because the island platform around St. Croix is much smaller and supports a smaller resident lobster population.

In Puerto Rico, divers have accounted for more lobster landings in recent years. Landings statistics indicate that divers accounted for 47,000 lbs (13% of total trap landings) in 1977 and 48,000 lbs (12%) in 1978 (hand and speared lobster; Weiler and Suarez-Caabro, 1980). A decade later divers accounted for more lobster and a greater percentage of the total trap landings: 65,222 lbs (83%) in 1988 and 53,232 lbs (42%) in 1989 (skin and SCUBA divers; Matos and Sadovy, 1990) [Note, Hurricane Allen may have affected 1989 landings and effort]. More information is needed about divers, particularly where they fish and the size-frequency of their landings.

Total annual landings averaged approximately half (53%) of the maximum sustained yield (MSY) estimated in the FMP (829,300 lbs per year). The reasons for the difference are unknown but are most likely due to any, or all of the following: overly optimistic MSY projections in the FMP, incomplete reporting of actual landings, and loss of yield due to landings of undersized lobster. As discussed later, the last factor is very likely to be important although its exact impact could not be quantified.

### Size-Frequency

Historical size-frequency data, where available, are shown for Puerto Rico (Table 2), St. Thomas/St. John (Table 3), and St. Croix (Table 4). Mean carapace length has remained fairly constant above 4 inches in the Virgin Islands but has declined in Puerto Rico from 4.4 inches in 1951 to 3.5 inches in 1989.

Length-frequency based on carapace lengths of sampled lobster were examined by sex classification for St. Croix, St. Thomas/St. John, and Puerto Rico (Table 5). Sex classifications were male, female without tar spots (spermatophores) or eggs, females with tar spots, and females with eggs. A few lobster, labeled in the data set as unidentified females, were not included in a specific sex classification, but were retained in the total length-frequency distribution. For comparative purposes these data were expressed in percent (Table 6) and cumulative percentages (Table 7). Puerto Rico lobster carapace lengths showed an approximately normal distribution around the minimum legal size of 3.5 in (89 mm) while both Virgin Islands locations showed a distinct absence of lobster below the minimum legal size (Tables 6 and 7).

Differences between coasts of Puerto Rico were examined using 1985 data for the south, west, and combined north and east coasts (Fig. 3). The latter were combined because of few existing data. Length-frequency patterns were generally consistent between coasts although there was a trend for the largest lobster to come from the

combined north and east coasts. The most likely explanation for these larger lobster is that some were probably caught further east closer to the Virgin Islands which tends to have larger lobster as discussed previously.

Size-frequency data were compared to a heavily fished spiny lobster fishery in Florida and unfished areas in the Dry Tortugas (Fig. 4) using data provided by Gregory et al., (1982) and Davis (1975). Lobster from all areas of the Virgin Islands and Puerto Rico tended to be larger than what was observed from Florida. Lobster from St. Croix and Puerto Rico tended to be smaller than those from the unfished Dry Tortugas. St. Thomas/St. John had a higher frequency of large lobster than Puerto Rico.

Size-frequency distributions were examined as a function of distance from shore in Puerto Rico in order to test the hypothesis that smaller lobster tended to be found closer to shore in shallow water, as in Florida. Distances examined were 0 - 3 nautical miles (n = 113 interviews), 3 - 6 nm (n = 87 interviews), and greater than 6 nm (n = 294). No significant difference in size-frequencies were noted with distance from shore (Figure 5). However, distance from shore did not necessarily reflect depth because the narrow shelf along the north and south coasts of Puerto Rico and the presence of islands to the east and west which were far offshore from fishing ports but still had shallow, "nearshore" water. Data on depth of capture were not available at the workshop.

### Minimum Size Compliance

The review team interpreted the absence of smaller lobster in Virgin Island catches as an indication of compliance with minimum legal size limits. Undersized lobster represented 40% of the total lobster landed from 1985 through 1989 from Puerto Rico. In contrast, the St. Croix data showed that undersized lobster represented 1.3% of the total lobster landed from 1987 through 1989. St. Thomas and St. John had only 2.9% undersized lobster for 1985 through 1989. Clearly minimum size limits were much more likely to be complied with in the Virgin Islands than in Puerto Rico based on size-frequency of landings.

Growth overfishing thus appears to be a major problem in Puerto Rico, based on the large number of undersized lobster being landed and the recent declines in total landings. A yield-per-recruit analysis would help quantify this situation, however, the review team, after considerable effort, was unable to generate an acceptable model because of a lack of growth data specifically tuned to Puerto Rico and the Virgin Islands (discussed later). Lyons and Kennedy (1980) found that harvesting of large numbers of small lobster could result in 68-83% loss to the fishery in Florida.

A model in the Lobster FMP (CFMC, pg 38) predicted effects of minimum size regulations on total landings. The model was calibrated to begin in 1980 and predictions were consistent with

actually observed patterns assuming that 3.5" minimum carapace size regulations were observed in the Virgin Islands and that status quo (no size limits) were being observed in Puerto Rico. Note, that a 3.5" carapace length was in effect within the Virgin Islands during this time before the Federal FMP went into effect in 1985. Declining total landings were predicted under the status quo (no size limits) which appears to be the situation in Puerto Rico (Fig. 6a), although total landings declined at a somewhat faster rate than predicted. Total landings were expected to remain relatively stable and perhaps increase somewhat with a 3.5 in minimum carapace size regulation which is consistent with what was observed in the Virgin Islands (Fig. 6b).

#### Catch-Per-Unit-Effort

A general consensus existed at the workshop that fishing effort has probably increased slowly over recent years. Although some data are available on the total number of fishermen (Table 1), effort data specifically targeting lobster were generally unavailable except some data for St. Croix. One problem is that lobster are caught by a variety of techniques including fish traps (pots), lobster traps (pots), and divers among others (Matos and Sadovy, 1990). In Puerto Rico, reported CPUE of lobsters landed (lbs/trap/yr) by fish traps, 34.3 (1977) and 29.2 (1978), was greater than by lobster traps, 24.1 (1977) and 15.1 (1978) (calculated from figures in Weiler and Caabro, 1980). The percentage of total lobster landed by lobster traps relative to

fish traps was small: 12% in 1977 and 9% in 1978 (Weiler and Caabro, 1980), and 24% in 1988 and 14% in 1989 (Matos and Sadovy, 1990).

Although a considerable amount of catch and effort data on a trip basis existed for Puerto Rico on the NMFS B6800 system in Miami, it was not considered useful for catch by trip analysis because there was no way to distinguish between completely and partially sampled trips. Although lobster are routinely caught by lobster and fish traps, it was not possible to distinguish from the data which trap type caught the lobster.

Analysis of catch-per-unit-effort (CPUE) for St. Croix from 1987 through 1989 based on monthly estimates of kilograms per trip and kilograms per pot (Fig. 7) show higher CPUE's in the winter and spring than in the summer and fall. Although Figure 7 also suggest that CPUE's may have declined over the 36 month sample period, not much confidence should be placed on a declining trend because data from only three months were available for 1987, these data came from winter months which tend to be high, and they are at one end of the regression series which gives them undue weight. More data over a longer period is necessary to define trends in CPUE with any confidence.

## Recommendations

The assessment team concluded that most obvious management action to increase the productivity of the spiny lobster fishery would be to increase compliance with minimum size restrictions in Puerto Rico. Spiny lobster growth studies are needed for Puerto Rico and the Virgin Islands to produce yield-per-recruit models. Studies should be directed describing the expanding diver-based spiny lobster fishery, particularly in Puerto Rico. Better data are needed on effort directed at spiny lobster and comparisons should be made of catch rates of spiny lobster in fish traps versus lobster traps. These two trap types will very likely have quite different catch efficiencies. Additional raw data from St. Croix on length-frequencies and catch-per-unit-effort should be entered in the database.

## Biological Parameters

### Growth

Determining growth is complex but important for properly managing the fishery (Hunt and Lyons, 1986). The assessment team concluded that insufficient data existed to properly characterize spiny lobster growth for Puerto Rico and the Virgin Islands region. It was agreed that growth parameters used in the spiny lobster FMP were probably unreliable having been based on early studies from the Virgin Islands in which Olsen et al. (1971) had reported a growth coefficient (K) equaled 0.43 for males and 0.32 for females. Munro (1983) estimated K as 0.21 when  $L_{\infty} = 190$  mm CL for Jamaica.

Estimates of spiny lobster growth coefficients range from 0.10 per year to 0.44 per year (Gulf of Mexico Fishery Management Council, 1982). Davis and Dodrill (1979) reported mean annual growth rates of 21.3 and 40.0 mm CL in Biscayne Bay and Florida Bay, respectively. Florida growth parameters, although well documented, were not considered appropriate because growth rates were likely to differ greatly due to different prevailing temperatures and stock conditions.

Considerable time was spent at the workshop attempting to estimate growth parameters for spiny lobster using the ELEFAN program (Pauly, 1985). The best available monthly length-frequency data to estimate growth were from St. Croix. Attempts to estimate growth parameters failed however for several reasons. First, carapace measurements were to the nearest one tenth inch which was too wide an interval to show distinct size-frequency peaks. Second, data were limited. Third, data were not available from individuals below the minimum size limit. Also, some assumptions of the ELEFAN program were violated because lobster grow in increments and lobster recruit throughout the year. CODREMAR had some growth data from very small tagged lobster but at too young an age to be useful.

After the workshop, a new study was found that examined spiny lobster growth in Jamaica. Haughton and Shaul (1989) gave a "first approximation" of spiny lobster growth for Jamaica at  $K = 0.48$  per

year and  $L_{\infty} = 193$  mm CL for males and  $K = 0.48$  per year and  $L_{\infty} = 193$  mm CL for females. These estimates were considered inadequate to use for Puerto Rico and the Virgin Islands because of lack of precision in the estimates and possible differences in stocks between areas (Haughton and Shaul, 1989), and concerns about the inappropriate application of the ELEFAN I program to lobster as discussed above. Without reliable growth parameters, yield-per-recruit models could not be generated.

### Mortality

It was not possible to estimate natural mortality from available data. Annual mortality was assumed to be 34% (equivalent to  $M = 0.42/\text{yr}$ ) in accordance with published literature from other locations (Waugh, 1981, Lyons and Hunt, 1987, Powers and Sutherland, 1989).

### Fecundity

Potential annual egg production was examined for Puerto Rico (Fig. 8), St. Croix (Fig. 9), and St. Thomas/St. John (Fig. 10) based on female size. Potential egg production assumes that each female reproduces only once and all females breed. These assumptions are unrealistic because not all females necessarily breed, especially smaller individuals (Lyons, et al. 1981) and some size classes may breed more than once per year. Potential egg production as illustrated probably overestimates relative egg contributions of smaller size classes while underestimating

contributions of larger size classes. Nevertheless these figures emphasize the importance of larger size classes to total egg production.

The Spawning Potential Ratio (SPR), the ratio of eggs produced between a fished and unfished population, was calculated from fishery dependent data according to methods used by Gregory, et al. (1982, his Tables 4 and 5) with available data from the most recent year for Puerto Rico (Table 8), St. Croix (Table 9) and St. Thomas (Table 10). Spawning potential was based on total mean fecundity, defined as the total number of eggs potentially produced divided by the total number of females (see Table 5 in Gregory, et al., 1982). Number of eggs per female was calculated according to the formula:

$$\text{Number of eggs} = 4.8(0.98 + 0.2598 \text{ CL})^{3.53},$$

where CL is carapace length in mm. Breeding females were considered females with spermatophores (tar spots) or eggs. The estimated total numbers of breeding females may be low because of legal prohibitions against landing egg bearing females (berried females). Attempts to calculate an Index of Reproductive Potential (Lyons, et al., 1981, their Fig. 13) failed because the results could not be calibrated with earlier studies; the 76-85 mm size class, used to calibrate curves, did not exist in Virgin Islands data.

Spawning potential, based on mean total fecundity, was compared to an unfished population in the Dry Tortugas and a heavily fished Florida population. For comparative purposes, 10 mm carapace length categories were used in calculations. However, calculations based on the midpoint of the carapace length provide some bias because the number of eggs increases exponentially with size. Therefore, calculations were also reported using 5 mm size categories and 1 mm size categories (see Tables 8 - 10).

Spawning potentials of 55.9% were calculated for Puerto Rico in comparison to an unfished population in the Dry Tortugas using 10 mm carapace length categories (see Gregory, 1982). This spawning potential is much higher than the 18.2% calculated for the Florida Keys for 1976 (Table 5 in Gregory, et al., 1982) or the 6% estimated for 1988 (GMFMC, Lobster Plan, Draft Amendment 3).

Calculated spawning potentials for the Virgin Islands exceeded the unfished Dry Tortugas population: 142% for St. Croix and 197% for St. Thomas. Although fundamental biological differences may exist between spiny lobster populations in the Virgin Islands and the Dry Tortugas, most of the difference can be explained as an artifact of the calculations. The Dry Tortugas estimate was based on actual catch from fishery independent sampling while the Virgin Islands estimates (and Puerto Rico) were based on commercial landings (fishery dependent) in which undersized individuals were excluded. Thus, very few females under 3.5" carapace length were

included in Virgin Islands data. This inflates mean total fecundity estimates because of the absence of numerous small, less fecund individuals in the calculations. Lyons et al. (1981) attempted to overcome this problem by standardizing data using a 76-85 mm carapace length as a basis for comparison. Unfortunately, this size category is one missing from Virgin Islands landings. A fishery independent sampling program would be necessary to better sample smaller size classes.

### Sex Ratios

Sex ratios (Males: Females) from available data since 1987 averaged 1.0 for Puerto Rico (Table 2), 1.6 for St. Thomas (Table 3), and 1.2 for St. Croix (Table 4). Sex ratios were skewed toward males in the Virgin Islands most likely because of larger lobster in the landings; males tend to get larger than females.

## CONCLUSIONS

### Status of Stocks

The spiny lobster fishery in the Virgin Islands appears healthy at present levels of fishing effort and under currently used fishing practices based on available data. Landings have remained consistent and the spawning potential appears high.

The spiny lobster assessment workshop panel viewed with particular alarm the nine-year decline in total landings and the

large number of undersized lobster being landed in Puerto Rico. Growth overfishing<sup>2</sup> appears to be a significant problem in Puerto Rico based on these facts. Recruitment overfishing<sup>3</sup> does not appear to be a problem under present levels of fishing effort based on calculated levels of spawning potential. The most reasonable explanation for these observations is that shallow water areas are being heavily exploited and overfished while deeper waters are less effectively exploited and maintain a reasonable number of large spawning individuals, some of which enter the landings (NOTE, the fact that no difference in size-frequency distributions were found with distance from shore does not refute this hypothesis). Thus, spawning potential appears high even though total landings are down. This scenario should be interpreted as a need to reduce fishing mortality on smaller lobster and not as an excuse to increase fishing effort on larger lobster in deeper water.

The assessment team concluded that most obvious management action to increase the productivity of the spiny lobster fishery would be to increase compliance with minimum size restrictions in Puerto Rico. Compliance appeared acceptable in the Virgin Islands.

---

<sup>2</sup> Growth overfishing occurs when fishes are caught too small, before they have had a chance to grow.

<sup>3</sup> Recruitment overfishing is a more serious problem that occurs when fishing reduces adult stocks such that lower egg production increases the chance of stock collapse recruitment failure.

The workshop did not deal with other potential issues including slot-size regulations, mortality caused by using undersized lobster used as bait in traps, degradable escape panels, or trap escape gaps which have been treated elsewhere (e.g. Lyons and Hunt, 1987; Powers and Sutherland, 1989). Although the original FMP discussed differences in landings between territorial and Exclusive Economic Zone (EEZ) waters, these could not be examined at the workshop because data that distinguished catch by location within or outside of the EEZ were unavailable.

#### Data Collection, Entry, and Management

Results of this workshop emphasize the continued need for standardized data collection, entry, and storage. Some analyses were hampered or were impossible because data were unavailable or stored in different formats.

#### Definition of Overfishing

The assessment panel was asked to comment on a definition of overfishing. Compared to Florida, the Virgin Islands and Puerto Rico show good representation of larger individuals which was interpreted to indicate lower fishing effort exists in both areas compared to the Florida spiny lobster fishery. The calculated spawning potential ratios were well above the 20% level recommended for a definition of overfishing by the Science and Statistical Committee. The 20% SPR was recommended based on theoretical

grounds (i.e. Goodyear 1989) and not on empirically derived stock-recruitment relationships which are unavailable for lobster. The lobster assessment workshop endorses the 20% SPR definition of overfishing. The 6% SPR recently proposed for Florida by the Gulf of Mexico Fishery Management Council Lobster Plan was based on a relatively long time period of empirical landings observations which are unavailable for the Caribbean region. The workshop participants considered it irresponsible to assume that the Caribbean region will respond to fishing pressure in the same way as southern Florida.

A definition of overfishing based solely on spawning potential appears to be inadequate, particularly considering the fact that total landings in Puerto Rico have declined for 9 years and are only 28 to 36% of peak values. One alternative is to include in the definition of overfishing a defined level of spawning potential and total landings. Ideally, the amount of total landings should be a percentage of some long-term average. It use is easier to define a level when landings have remained relatively stable such as in the Virgin Islands or in southern Florida (Powers and Sutherland, 1989). In Puerto Rico, however, there was no period of stable landings existing to use as a baseline. A possible definition submitted for Council consideration is:

"A spiny lobster stock is considered overfished when any of the following are observed: the spawning potential ratio is less than 20%, when total landings have declined to a level below 75% of the 5-year running mean, or when total landings have declined for 3 consecutive years."

With this definition, the Puerto Rico fishery became overfished in 1983 when landings dropped below 318,000 lbs and remained overfished until 1989 when landings increased from 143,761 to 186,423 lbs (Figure 11). Unfortunately, the 1989 levels are still well below those in previous years (Table 1) although technically they are not overfished by this definition. One way to deal with this problem would be to include a definition stating that:

"When overfished a stock will continue to be considered overfished until the SPR is above 20% and total landings are above the level at which the fishery first became overfished" (i.e. 380,000 lbs).

Obviously other levels of landings could be considered.

#### SUMMARY OF MAJOR RECOMMENDATIONS

1. Data collection, entry, and storage should be standardized as much as possible.
2. Where possible, data entry should be done by data collecting entities to avoid misinterpretation.
3. Raw data from St. Croix on length-frequencies and catch-per-unit-effort should be entered in the database.
4. Compliance with minimum sizes and other regulations should be increased, particularly in the Puerto Rico fishery.
5. Growth and mortality studies are needed for Puerto Rico and the Virgin Islands to produce yield-per-recruit models.
6. The diver-based spiny lobster fishery in Puerto Rico should be studied in terms of total effort, areas fished, and size composition of landings.

7. Better fishing effort data are needed.
8. Comparisons should be made of catch rates of spiny lobster in fish traps versus lobster traps.
9. Fishery independent sampling of lobster size-frequency distributions are needed to better estimate spawning potential.
10. A modified definition of overfishing is recommended that considers total landings as well as spawning potential.

#### ACKNOWLEDGEMENTS

We thank John Hunt (Florida DNR, Marathon, FL) and Nancy Thompson (NMFS, Miami Laboratory) for critical comments on the draft report. Ausbon Brown Jr. coordinated data entry. Doug Harper and David McClellan (NMFS, SEFC, Miami) assisted with analyses.

#### LITERATURE CITED

- CFMC. 1981. Environmental impact statement, fishery management plan and regulatory impact review for the spiny lobster fishery of Puerto Rico and the U.S. Virgin Islands. Caribbean Fishery Management Council. 66 p.
- CFMC. 1982. Report on the spiny lobster size frequency survey 1980-81. Caribbean Fishery Management Council. 22 p.
- Davis, G.E. 1975. Minimum size of mature spiny lobster, Panulirus argus at Dry Tortugas, Florida. Trans. Am. Fish. Soc. 104: 675-676.
- Feliciano, C. 1958. The lobster fishery of Puerto Rico. Proc. Gulf and Carib. Fish. Inst. 10: 147-156.
- Goodyear, C.P. 1989. Spawning stock biomass per recruit: The biological basis for a fisheries management tool. ICCAT Working Document SCRS/89/82.

- Gregory, D.R., Jr., R.F. Labisky, and C.L. Combs. 1982. Reproductive dynamics of the spiny lobster Panulirus argus in South Florida. Transactions of the American Fisheries Society 111: 575-584.
- Gulf of Mexico Fishery Management Council. 1982. Fishery Management Plan, Environmental Impact Statement and Regulatory Impact Review for spiny lobster in the Gulf of Mexico and South Atlantic, SAFMC, 1 Southpark Circle, Suite 306, Charleston, SC, 29407-4699.
- Haughton, M. and W. Shaul. 1989. Estimation of growth parameters for the spiny lobster (Panulirus argus) in Jamaican Waters. Proc. Gulf and Carib. Fish. Inst. 39: 279-288.
- Hunt, J.H. and W.G. Lyons. 1986. Factors affecting growth and maturation of spiny lobsters, Panulirus argus, in the Florida Keys. Can. J. Fish. Aquat. Sci. 43: 2243-2247.
- Lyons, W.G., D.G. Barber, S.M. Foster, F.S. Kennedy, Jr., and G.R. Milano. 1981. The spiny lobster, Panulirus argus, in the Middle and Upper Florida Keys: Population structure, seasonal dynamics, and reproduction.
- Lyons, W.G. and Hunt, J.H. 1987. Catch rates of spiny lobsters, Panulirus argus, in traps equipped with escape gaps, and potential benefits to the south Florida fishery. Proc. Gulf Carib. Fish. Inst. 39 (in press).
- Lyons, W.G. and F.S. Kennedy, Jr. 1980. Effects of harvest techniques on sublegal spiny lobsters and on subsequent fishery yield. Proc. Gulf Carib. Fish. Inst. 33: 290-300.
- Matos, D. and Y. Sadovy. 1990. Overview of Puerto Rico's small-scale fisheries statistics (Perspectivas de las Estadísticas de la Pesca en Pequeña Escala de Puerto Rico) 1988-1989. Technical Report CODREMAR 1(4): 1-17.
- Matos, D. and Y. Sadovy. 1990b. CODREMAR/NMFS Cooperative Statistics Program for 1989-1990. Annual Report to the National Marine Fisheries Service. Project SF-33. June 1990. 44 p.
- Mattox, N.T. 1952. A preliminary report on the biology and economics of the spiny lobster in Puerto Rico. Proc. Gulf Carib. Fish. Inst. 4: 69-70.
- Munro, J.L. 1983. The biology, ecology and bionomics of spiny lobsters (Palinuridae), spider crabs (Majidae) and other crustacean resources. Pages In: J.L. Munro (ed). Caribbean coral reef fishery resources. International Center for Living Aquatic Resources Management (ICLARM). Manila, Philippines.

- Olsen, D.A., W.F. Hernkind and R.A. Cooper. 1975. Population dynamics, ecology and behavior of spiny lobsters, Panulirus argus, of St. John, U.S.V.I.: I, Introduction and general population characteristics. In S.A. Earle and R.J. Lavenberg (ed.). Results of the Tektite program: Coral reef invertebrates and plants. Nat. Hist. Mus. Los Ang. Cty. Sci. Bull 20: 11-16.
- Olsen, D.A. and I.G. Kobic. 1975. Population dynamics, ecology and behavior of spiny lobsters, Panulirus argus, of St. John, U.S.V.I.: II Growth and mortality. In S.A. Earle and R.J. Lavenberg (ed.). Results of the Tektite program: Coral reef invertebrates and plants. Nat. Hist. Mus. Los Ang. Cty. Sci. Bull 20: 17-21.
- Pauly, D. 1985. The population dynamics of short-lived species, with emphasis on squids. NAFO Scientific Council Studies. No. 9: 143-154.
- Powers, J.E. and D.L. Sutherland. 1989. Spiny lobster assessment, CPUE, size-frequency, yield per recruit, and escape gap analyses. Miami Laboratory, Southeast Fisheries Center, National Marine Fisheries Service, NOAA, Coastal Resources Division Contribution Number CRD-88\89-24.
- Waugh, G.T. 1981. Management of juvenile spiny lobster (Panulirus argus) based on estimated biological parameters from Grand Bahama Island, Bahamas. Proc. Gulf Carib. Fish Inst. 33: 271-289.
- Weiler, D. and J. Suarez-Caabro. 1980. Overview of Puerto Rico's small-scale fisheries statistics (Perspectivas de las Estadisticas de la Pesca en Pequena Escala de Puerto Rico) 1972-1978. Informe Tecnico, CODREMAR 1(1): 1-17.

## List of Tables

- Table 1. Total Landings and Effort Summary.
- Table 2. Size-Frequency Surveys of Spiny Lobster for Puerto Rico.
- Table 3. Size-Frequency Surveys of Spiny Lobster for St. Thomas and St. John.
- Table 4. Size-Frequency Surveys of Spiny Lobster for St Croix.
- Table 5. Length-frequencies for spiny lobsters.
- Table 6. Percent length-frequencies for spiny lobsters.
- Table 7. Cumulative length-frequencies for spiny lobsters.
- Table 8. Fecundity Estimates for Spiny Lobster from Puerto Rico.
- Table 9. Fecundity Estimates for Spiny Lobster from St. Croix.
- Table 10. Fecundity Estimates for Spiny Lobster from St. Thomas/St. John.

## List of Figures

- Figure 1. Total Lobster Landings for Puerto Rico.
- Figure 2. Total Lobster Landings for the Virgin Islands.
- Figure 3. Female length-frequency and potential egg production based on carapace lengths for different coasts of Puerto Rico.
- Figure 4. Female length-frequency and potential egg production based on carapace lengths for Florida locations.
- Figure 5. Lobster length-frequencies with distance from Puerto Rico.
- Figure 6. Predicted and observed total lobster landings for the Virgin Islands and Puerto Rico. Predicted trends in the Lobster FMP were calibrated to start with actual reported 1981 landings. (Top) Predicted Virgin Islands landings were based on a 3 in minimum carapace length (effective in VI) until 1985 when a 3.5 in carapace length became effective. (Bottom) Predicted Puerto Rico landings were based on the status quo model (no regulations) in the Lobster FMP.
- Figure 7. Catch-per-unit-effort for St. Croix (1987-1989). (Top) CPUE per trip. (Bottom) CPUE per pot haul.
- Figure 8. Female length-frequency and potential egg production based on carapace lengths for Puerto Rico.
- Figure 9. Female length-frequency and potential egg production based on carapace lengths for St. Croix.
- Figure 10. Female length-frequency and potential egg production based on carapace lengths for St. Thomas/St. John.
- Figure 11. Total landings compared to 75% of a 5-year running mean. The running mean was calculated by including the previous four years of landings with the current year's landings.

### List of Participants

Dr. Richard Appeldoorn, Department of Marine Sciences, University of Puerto Rico, Mayaguez, Puerto Rico 00708.


Jim Beets, Division of Fish and Wildlife, Department of Conservation and Cultural Affairs, 101 Estate Nazareth, St. Thomas, VI 00802.

Dr. James A. Bohnsack, National Marine Fisheries Service, Southeast Fisheries Center, Miami Laboratory, 75 Virginia Beach Dr., Miami, Florida 33149.


Steve Meyers, Caribbean Fishery Management Council, Suite 1108 Banco de Ponce Bldg., Hato Rey, Puerto Rico 00918

Dr. Yvonne Sadovy, Department of Natural Resources, P.O. Box 3665, Marina Station, Mayaguez, Puerto Rico 00709.


# Puerto Rico Total Lobster Landings




# Virgin Islands Total Lobster Landings




Puerto Rico, 1984 North & East Coast  
(n = 112)



Puerto Rico, 1985 South Coast  
(n = 219)




Puerto Rico, 1985 West Coast  
(n = 214)




57


**Dry Tortugas, 1973-75**  
(n = 1594)






**Florida Keys, 1976**  
(n = 782)



**Florida Keys, 1978-79**  
(n = 1594)




# LOBSTER LENGTH FREQUENCY FROM PUERTO RICO




## Puerto Rico Total Lobster Landings

Observed versus Predicted




## Virgin Islands Lobster Landings


Observed versus Predicted










## Spiny Lobster 1990 Stock Assessment




## Spiny Lobster 1990 Stock Assessment







St. Croix, 1970  
(n = 67)




St. Croix, 1988  
(n = 438)




St. Croix, 1987  
(n = 297)




St. Croix, 1989  
(n = 245)




St. Thomas, 1984  
(n = 99)




St. Thomas, 1986  
(n = 468)




St. Thomas, 1985  
(n = 481)




St. Thomas, 1987  
(n = 167)




St. Johns, 1985  
(n = 787)



St. Thomas, 1988  
(n = 115)



# Puerto Rico Total Lobster Landings



### List of Tables

- Table 1. Total Landings and Effort Summary.
- Table 2. Size-Frequency Surveys of Spiny Lobster for Puerto Rico.
- Table 3. Size-Frequency Surveys of Spiny Lobster for St. Thomas and St. John.
- Table 4. Size-Frequency Surveys of Spiny Lobster for St Croix.
- Table 5. Length-frequencies for spiny lobsters.
- Table 6. Percent length-frequencies for spiny lobsters.
- Table 7. Cumulative length-frequencies for spiny lobsters.
- Table 8. Fecundity Estimates for Spiny Lobster from Puerto Rico.
- Table 9. Fecundity Estimates for Spiny Lobster from St. Croix.
- Table 10. Fecundity Estimates for Spiny Lobster from St. Thomas/St. John.

Table 1. Total Landings and Effort Summary.

Puerto Rico			St Thomas and St. Johns			St Croix					
Year	Total Landings	Total Fishers	Total Vessels	Year	Total Landings	Licensed Fishers	Total Vessels	Year	Total Landings	Licensed Fishers	Total Vessels
1951	466760		223								
1964	150000										
1969	354000										
1970	417000										
1971	258000										
1972	237000		970								
1973	250000		930								
1974	244000		1120								
1975	311000		1230				865				
1976	384000		1230				901				
1977	421000		1368				1036				
1978	451000		1442				1073				
1979	512000		1442				1073				
1980	474000		1447				1087				
1981	481000			80-81	29418		258		7148		163
1982	359000	1872	1449	81-82	47204		256		8280		322
1983	294229	1415	1125	82-83	29460		259		2304		195
1984	283262			83-84	39810		255		7419		182
1985	246501	1766		84-85	41911		255		8328		182
1986	219203	1135	865	85-86	39300		330		16031		206
1987	158223	1731		86-87	23296		329		4322		200
1988	143761			87-88	41875		306		4437		217
1989	186423	1822	1107	88-89							
Mean	317451	1395	1058		36534		281		7284		208

Table 2. Size-Frequency Surveys of Spiny Lobster for Puerto Rico.

Survey	Year	Number of Lobster Sampled	Mean Carapace Length (in)	Mean Carapace Length (mm)	Mean Weight (lbs)	Percent Below 3.5 in (Numbers)	Percent Below 3.5 in (lbs)	Sex			Max. Carapac Length (mm)	Mean Weight (g)
								Females	Males	M:F		
Mattox, 1952	1951		4.4	113								
Feliciano, C.	1956-5	1276	4.0	101.6	2.0	19.6	-					
CFMC, 1981	1968	223	3.8	95.3	1.7	25.0	-					
Olsen & Koblic	1975		4.3	109.3		15.1						
CFMC, 1981	1978-7	9232	3.7	93.5	1.7	40.5	23.7					
CODREMAR*	1980	129	3.7	92.8		27.0		75	54	0.72	127	
CFMC, 1982	1980-1	5574	3.8	95.3	1.8	34.7						
CODREMAR	1982											
CODREMAR	1983	211	3.7	94.4		28.0		106	105	0.990	152	
CODREMAR	1984	2184				31.0		1093	1091	0.998		
CODREMAR (all)	1985					32.0						
" South Coast	"	554						297	257	0.865		
" N & E Coast	"	271						135	136	1.007	168	
" West Coast	"	480						235	245	1.042	163	
CODREMAR	1986	568	3.6	92.5		39.0		258	310	1.201	174	
CODREMAR	1987	387	3.8	95.6		30.0		179	208	1.162	152	
CODREMAR	1988	52						31	21	0.677		
CODREMAR	1989	392	3.5	90.1		41.0		235	276	1.174		
Matos & Sadovy, 1990b	1989	1037	3.5	90								

\* Data collected by CODREMAR and available to the workshop on the NMFS TIP database.

Table 3. Size-Frequency Surveys of Spiny Lobster for St Thomas and StSt Thomas and St Johns, U.S. Virgin Islands

Survey	Year	Number of Lobster Sampled	Mean Carapace Length (in)	Mean Carapace Length (mm)	Mean Weight (lbs)	Percent Below 3.5 in (Numbers)	Percent Below 3.5 in (lbs)	Sex			Max. Carapac Length (mm)	Mean Weight (g)
								Females	Males	M:F		
St. John												
DCCA USVI*	1985	1802	4.1	105				790	1012	1.281	152	
St. Thomas												
CFNC, 1981 +	1978	146	4.4	112	2.6	9.6	6.1					
CFNC, 1982	1979	89	4.4	113	2.8	7.9						
	1980											
CFNC, 1982	1981	89	4.5	114	2.8							
DCCA USVI*	1982	689	4.5	114		16.7	#					
DCCA USVI	1983	107	4.2	106								
DCCA USVI	1984	219	4.5	115	2.7	5.0		99	120	1.212	191	
DCCA USVI	1985	1060	4.6	116	2.6	0.7		481	564	1.172	203	
DCCA USVI	1986	1345	4.3	109	2.4	1.7		468	846	1.807	191	
DCCA USVI	1987	368	4.7	119	3.0	0.3		167	200	1.197	178	
DCCA USVI	1988	313	4.4	111	2.6	0.0		115	198	1.721	165	

\* Data collected by Dept. of Conservation and Community Affairs and available to the workshop on the NMFS TIP database.

+ June data only

Table 4. Size-Frequency Surveys of Spiny Lobster for St Croix, U.S.V.I.

Survey	Year	Number of Lobster	Mean	Mean	Mean Weight (lbs)	Percent	Percent	Max.			
			Carapace Length (in)	Carapace Length (mm)		Below 3.5 in (Numbers)	Below 3.5 in (lbs)	Sex Ratio M:F	Carapac Length (mm)	Mean Weight (g)	
Olsen et al., 1975	1970-1	756	4.4	113							
CFMC, 1981	1976	996	4.1	103	2.0	1.0	-				
	1977										
CFMC, 1981 #	1978	233	4.6	117	2.6	0.4	2.7				
CFMC, 1982	1979	90	4.3	109		15.5 *					
	1980										
CFMC, 1982	1981	90	4.3	109	2.5						
DCCA USVI +	1981		3.9	99							
DCCA USVI	1982	482	4.1	105		25.9 *					
DCCA USVI	1983	41	3.8	96							
DCCA USVI	1984	383	4.1	104							
	1985										
	1986										
DCCA USVI	1987	637	4.1	105	2.2	2.7	297	340	1.144	150	989
DCCA USVI	1988	965	4.2	106	2.1	1.3	438	522	1.191	150	976
DCCA USVI	1989	578	4.2	106	2.2	1.4	245	333	1.359	152	983

\* includes "legal" lobster 3.5 and 3.6" CL.

+ Available data collected by Dept. of Conservation and Community Affairs on the NMFS TIP database.

# July data only

Table 5: Length-frequencies for spiny lobsters (*Panulirus argus*) for St. Croix (1987-1989), St. Thomas and St. John (1985-1989), and Puerto Rico (1985-1989). Note, columns will not add up exactly because a few lobsters labeled in the data set as unidentified females, were not included in a specific column based on sex type, but were retained in the total columns.

LGTH (IN)	ST. CROIX					ST. THOMAS AND ST. JOHN					PUERTO RICO				
	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)
2.8						6				6	354	50	4		288
2.9						12				12	669	69	8		588
3						43	1			42	741	96	36	7	594
3.1	1	1				40		4		36	721	89	88	14	520
3.2	2	2				24	2	4		18	745	96	40	7	594
3.3	15	3	8		4	133	11	32		90	798	120	160	14	498
3.4	35	7	16		12	253	25	72		156	881	118	240	7	510
3.5	217	34	96	7	80	765	89	216	7	450	977	107	272	42	552
3.6	319	56	176		66	562	83	244	14	216	790	114	308	56	312
3.7	361	68	220	21	46	609	107	276	28	198	703	131	280	42	246
3.8	338	63	232	7	36	560	113	300	42	102	723	99	340	28	252
3.9	337	55	236		34	580	122	304	28	126	897	161	424	42	258
4	352	75	236	7	28	728	162	344	21	198	626	104	300	28	192
4.1	329	88	184	7	44	526	132	244	42	108	614	171	312	21	108
4.2	344	94	224		26	569	153	284	21	108	323	96	148	7	72
4.3	322	68	196	14	32	645	150	308	28	156	471	112	224	35	96
4.4	232	73	136	7	16	427	137	184	7	96	328	80	136	28	84
4.5	178	70	100		8	593	145	284	14	144	251	67	88	42	54
4.6	172	60	96		16	322	107	136	28	48	152	47	56	7	42
4.7	114	36	68		10	223	80	88	7	48	125	26	44	7	48
4.8	110	42	60		8	306	93	180		30	128	46	52		30
4.9	81	43	36		2	175	44	64	7	60	55	24	12	7	12
5	43	23	16		4	311	111	136	28	36	96	25	24	21	24
5.1	38	22	12		4	78	40	32		6	39	15	16		6
5.2	37	13	20		4	181	76	84		18	55	19	24		12
5.3	16	10	4		2	57	43	8		6	27	15	12		
TOTAL	3993	1006	2372	70	482	8728	2026	3828	322	2514	12289	2097	3648	462	5992

Table 6: Percent length-frequencies for spiny lobsters (*Panulirus argus*) for St. Croix (1987-1989), St. Thomas and St. John (1985-1989), and Puerto Rico (1985-1989).

LGTH (IN)	ST. CROIX					ST. THOMAS AND ST. JOHN					PUERTO RICO				
	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)
2.8						0.07%					0.07%				
2.9						0.14%					0.14%				
3						0.49%	0.01%				0.48%				
3.1	0.03%	0.03%				0.46%		0.05%			0.41%				
3.2	0.05%	0.05%				0.27%	0.02%	0.05%			0.21%				
3.3	0.38%	0.08%	0.20%		0.10%	1.52%	0.13%	0.37%			1.04%				
3.4	0.88%	0.18%	0.41%		0.31%	2.90%	0.29%	0.83%			1.80%				
3.5	5.43%	0.87%	2.44%	0.18%	2.04%	8.76%	1.02%	2.49%	0.08%	5.18%	7.95%	0.88%	2.23%	0.34%	4.52%
3.6	7.99%	1.42%	4.48%		1.68%	6.44%	0.96%	2.81%	0.16%	2.49%	6.43%	0.93%	2.52%	0.46%	2.56%
3.7	9.04%	1.73%	5.60%	0.53%	1.17%	6.98%	1.23%	3.18%	0.32%	2.28%	5.72%	1.07%	2.30%	0.34%	2.02%
3.8	8.46%	1.60%	5.90%	0.18%	0.92%	6.42%	1.30%	3.45%	0.48%	1.17%	5.88%	0.81%	2.79%	0.23%	2.07%
3.9	8.44%	1.40%	6.01%		0.87%	6.65%	1.40%	3.50%	0.32%	1.45%	7.30%	1.32%	3.48%	0.34%	2.11%
4	8.82%	1.91%	6.01%	0.18%	0.71%	8.34%	1.86%	3.96%	0.24%	2.28%	5.09%	0.85%	2.46%	0.23%	1.57%
4.1	8.24%	2.24%	4.68%	0.18%	1.12%	6.03%	1.52%	2.81%	0.48%	1.24%	5.00%	1.40%	2.56%	0.17%	0.89%
4.2	8.62%	2.39%	5.70%		0.66%	6.52%	1.76%	3.27%	0.24%	1.24%	2.63%	0.79%	1.21%	0.06%	0.59%
4.3	8.06%	1.73%	4.99%	0.36%	0.81%	7.39%	1.73%	3.54%	0.32%	1.80%	3.83%	0.92%	1.84%	0.29%	0.79%
4.4	5.81%	1.86%	3.46%	0.18%	0.41%	4.89%	1.58%	2.12%	0.08%	1.10%	2.67%	0.66%	1.11%	0.23%	0.69%
4.5	4.46%	1.78%	2.54%		0.20%	6.79%	1.67%	3.27%	0.16%	1.66%	2.04%	0.55%	0.72%	0.34%	0.44%
4.6	4.31%	1.53%	2.44%		0.41%	3.69%	1.23%	1.57%	0.32%	0.55%	1.24%	0.39%	0.46%	0.06%	0.34%
4.7	2.85%	0.92%	1.73%		0.25%	2.55%	0.92%	1.01%	0.08%	0.55%	1.02%	0.21%	0.36%	0.06%	0.39%
4.8	2.75%	1.07%	1.53%		0.20%	3.51%	1.07%	2.07%		0.35%	1.04%	0.38%	0.43%		0.25%
4.9	2.03%	1.09%	0.92%		0.05%	2.01%	0.51%	0.74%	0.08%	0.69%	0.45%	0.20%	0.10%	0.06%	0.10%
5	1.08%	0.59%	0.41%		0.10%	3.56%	1.28%	1.57%	0.32%	0.41%	0.78%	0.20%	0.20%	0.17%	0.20%
5.1	0.95%	0.56%	0.31%		0.10%	0.89%	0.46%	0.37%		0.07%	0.32%	0.12%	0.13%		0.05%
5.2	0.93%	0.33%	0.51%		0.10%	2.07%	0.87%	0.97%		0.21%	0.45%	0.16%	0.20%		0.10%
5.3	0.40%	0.25%	0.10%		0.05%	0.65%	0.49%	0.09%		0.07%	0.22%	0.12%	0.10%		
TOTAL	100.00%	25.60%	60.36%	1.78%	12.26%	100.00%	23.31%	44.05%	3.71%	28.93%	100.00%	17.19%	29.90%	3.79%	49.12%

Table 7: Cumulative percent length-frequencies for spiny lobsters (*Panulirus argus*) for St. Croix (1987-1989), St. Thomas and St. John (1985-1989), and Puerto Rico (1985-1989).

LGTH (IN)	ST. CROIX					ST. THOMAS AND ST. JOHN					PUERTO RICO				
	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)	ALL LOBSTER	MALE	FEMALE (TAR)	FEMALE (EGGS)	FEMALE (OTHER)
2.8						0%				0%	3%	2%	0%		5%
2.9						0%				1%	8%	6%	0%		15%
3						1%	0%			2%	14%	10%	1%	2%	25%
3.1	0%	0%				1%	0%	0%		4%	20%	14%	4%	5%	33%
3.2	0%	0%				1%	0%	0%		5%	26%	19%	5%	6%	43%
3.3	0%	1%	0%		1%	3%	1%	1%		8%	33%	25%	9%	9%	51%
3.4	1%	1%	1%		3%	6%	2%	3%		14%	40%	30%	16%	11%	60%
3.5	7%	5%	5%	10%	20%	15%	6%	9%	2%	32%	48%	36%	23%	20%	69%
3.6	15%	10%	12%	10%	34%	21%	10%	15%	7%	41%	54%	41%	32%	32%	74%
3.7	24%	17%	22%	40%	43%	28%	16%	22%	15%	49%	60%	47%	39%	41%	78%
3.8	32%	23%	32%	50%	51%	34%	21%	30%	28%	53%	66%	52%	49%	47%	83%
3.9	41%	29%	41%	50%	58%	41%	27%	38%	37%	58%	73%	60%	60%	56%	87%
4	50%	36%	51%	60%	63%	49%	35%	47%	43%	66%	78%	65%	69%	62%	90%
4.1	58%	45%	59%	70%	73%	55%	42%	53%	57%	70%	83%	73%	77%	67%	92%
4.2	66%	54%	69%	70%	78%	62%	49%	61%	63%	74%	86%	77%	81%	68%	93%
4.3	74%	61%	77%	90%	85%	69%	57%	69%	72%	80%	90%	83%	87%	76%	95%
4.4	80%	68%	83%	100%	88%	74%	64%	74%	74%	84%	92%	86%	91%	82%	96%
4.5	85%	75%	87%		90%	81%	71%	81%	78%	90%	94%	90%	93%	91%	97%
4.6	89%	81%	91%		93%	85%	76%	85%	87%	92%	96%	92%	95%	92%	98%
4.7	92%	85%	94%		95%	87%	80%	87%	89%	94%	97%	93%	96%	94%	99%
4.8	95%	89%	96%		97%	91%	85%	92%	89%	95%	98%	95%	98%	94%	99%
4.9	97%	93%	98%		97%	93%	87%	93%	91%	97%	98%	96%	98%	95%	99%
5	98%	96%	98%		98%	96%	92%	97%	100%	99%	99%	98%	99%	100%	100%
5.1	99%	98%	99%		99%	97%	94%	98%		99%	99%	98%	99%		100%
5.2	100%	99%	100%		100%	99%	98%	100%		100%	100%	99%	100%		100%
5.3	100%	100%	100%		100%	100%	100%	100%		100%	100%	100%	100%		100%

Table 8. Fecundity Calculations for Puerto Rico (1989, West Coast).

Carapace Length (mm)	Percentage Contribution			Fecundity <sup>1</sup>			
	Number of Females	Number of Breeding Females	Annual Egg Production	Estimated Number of Eggs (x10 <sup>3</sup> )	Puerto Rico 1989 (x10 <sup>3</sup> )	Florida Keys 1976 (x10 <sup>3</sup> )	Tortugas 1973-75 (x10 <sup>3</sup> )
<65	0	0	0	0.0	0.0	0.0	0.0
65-75	19	3	1	323.3	7.5	21.7	0.0
75-85	29	11	6	2025.6	31.6	68.8	63.3
85-95	23	34	26	9424.9	184.8	113.1	225.8
95-105	18	32	34	12399.7	302.4	380.6	394.0
105-115	8	15	22	8205.7	455.9	746.0	552.5
115-125	2	4	8	3013.0	753.2	0.0	761.9
125-135	1	1	4	1321.3	660.6	0.0	681.9
135-145	0	0	0	0.0	0.0	0.0	0.0
Total	100	100	100	36713.4	2396.2	1330.2	2679.4
Sample Size	223	73	73				
Total Mean Fecundity (uncorrected)				3	164.6	-	-
Total Mean Fecundity (corrected, n = 78 breeders)					175.9	57.2	314.7
Total Mean Fecundity (calculated with 5 mm classes)					177.9	-	-
Total Mean Fecundity (calculated with 1 mm classes)					182.5	-	-
Spawning Potential Ratio (10 mm classes)					55.89%	18.18%	

NOTES:

1 Fecundity = (number of eggs)/(number of females).

2 Number of Eggs = Number of breeding females x  $[4.8(0.98 + 0.2598CL)^{3.53}]$   
where CL is the midpoint of the carapace-length class.

3 Corrected mean was calculated by multiplying the mean by 78/73 to account for 5 reproductive females without carapace measurements.

Table 9. Fecundity Calculations for St. Croix (1989).

Carapace Length (mm)	Percentage Contribution			Fecundity <sup>1</sup>			
	Number of Females	Number of Breeding Females	Annual Egg Production	Estimated Number of Eggs (x10 <sup>3</sup> )	St. Croix 1989 (x10 <sup>3</sup> )	Florida Keys 1976 (x10 <sup>3</sup> )	Tortugas 1973-75 (x10 <sup>3</sup> )
<65	0.0	0.0	0.0	0.0	0.0	0.0	0.0
65-75	0.0	0.0	0.0	0.0	0.0	21.7	0.0
75-85	0.8	0.6	0.2	253.2	126.6	68.8	63.3
85-95	20.8	13.7	7.9	8670.9	170.0	113.1	225.8
95-105	41.2	43.5	36.0	39355.5	389.7	380.6	394.0
105-115	24.9	26.8	30.7	33568.9	550.3	746.0	552.5
115-125	9.8	13.1	20.2	22095.2	920.6	0.0	761.9
125-135	2.4	2.4	4.8	5285.0	880.8	0.0	681.9
135-145	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total	100	100	100	109228.7	3038.059	1330.2	2679.4
n =	245	168	168				
Total Mean Fecundity (10 mm size classes)					445.8	57.2	314.7
Total Mean Fecundity (5 mm classes)					-	-	-
Total Mean Fecundity (0.1 in classes)					0.0	-	-
Spawning Potential Ratio					141.67%	18.18%	

## NOTES:

1 Fecundity = (number of eggs)/(number of females).

2 Number of Eggs = Number of breeding females x  $[4.8(0.98 + 0.2598CL)^{3.53}]$   
 where CL is the midpoint of the carapace-length class.

Table 10. Fecundity Calculations for St. Thomas (1988).

Carapace Length (mm)	Percentage Contribution			1 Fecundity			
	Number of Females	Number of Breeding Females	Annual Egg Productio	Estimated Number of Eggs (x10 <sup>3</sup> )	St. Croix 1989 (x10 <sup>3</sup> )	Florida Keys 1976 (x10 <sup>3</sup> )	Florida Tortugas 1973-75 (x10 <sup>3</sup> )
<65	0.0	0.0	0.0	0.0	0.0	0.0	0.0
65-75	0.0	0.0	0.0	0.0	0.0	21.7	0.0
75-85	0.0	0.0	0.0	0.0	0.0	68.8	63.3
85-95	31.6	27.1	14.2	8670.9	279.7	113.1	225.8
95-105	19.4	20.0	15.1	9165.0	482.4	380.6	394.0
105-115	27.6	30.6	31.9	19395.3	718.3	746.0	552.5
115-125	10.2	10.6	14.9	9039.0	903.9	0.0	761.9
125-135	10.2	10.6	19.5	11891.3	1189.1	0.0	681.9
135-145	0.0	0.0	0.0	0.0	0.0	0.0	0.0
145-155	0.0	0.0	0.0	0.0	0.0	0.0	0.0
155-165	1.0	1.2	4.4	2698.9	2698.9	0.0	0.0
Total	100	100	100	60860.44	6272.337	1330.2	2679.4
n =	98	85	85				
Total Mean Fecundity (10 mm size classes)					621.0	57.2	314.7
Total Mean Fecundity (5 mm classes)					-	-	-
Total Mean Fecundity (0.1 in classes)					583.8	-	-
Spawning Potential Ratio					197.34†	18.18†	

## NOTES:

1 Fecundity = (number of eggs)/(number of females).

2 Number of Eggs = Number of breeding females x [4.8(0.98 + 0.2598CL)<sup>3.53</sup>]  
where CL is the midpoint of the carapace-length class.

Figure 11

Puerto Rico Total Lobster Landings

